



Biodegradation: Updating the Concepts of Control for Microbial Cleanup in Contaminated Aquifers

Meckenstock, Rainer U.; Elsner, Martin; Griebler, Christian; Lueders, Tillmann; Stumpp, Christine; Aamand, Jens; Agathos, Spiros N.; Albrechtsen, Hans-Jørgen; Bastiaens, Leen; Bjerg, Poul Løgstrup

Total number of authors:
18

Published in:
Environmental Science and Technology

Link to article, DOI:
[10.1021/acs.est.5b00715](https://doi.org/10.1021/acs.est.5b00715)

Publication date:
2015

Document Version
Peer reviewed version

[Link back to DTU Orbit](#)

Citation (APA):

Meckenstock, R. U., Elsner, M., Griebler, C., Lueders, T., Stumpp, C., Aamand, J., Agathos, S. N., Albrechtsen, H.-J., Bastiaens, L., Bjerg, P. L., Boon, N., Dejonghe, W., Huang, W. E., Schmidt, S. I., Smolders, E., Sørensen, S. R., Springael, D., & van Breukelen, B. M. (2015). Biodegradation: Updating the Concepts of Control for Microbial Cleanup in Contaminated Aquifers. *Environmental Science and Technology*, 49(12), 7073-7081. <https://doi.org/10.1021/acs.est.5b00715>

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Feature

Biodegradation: Updating the concepts of control for microbial clean-up in contaminated aquifers

Rainer Udo Meckenstock, Martin Elsner, Christian Griebler, Tillmann Lueders, Christine Stumpp, Winnie Dejonghe, Leen Lucia Bastiaens, Dirk Springael, Erik Smolders, Nico Boon, Spiros N. Agathos, Sebastian R. Sorensen, Jens Aamand, Hans Jorgen Albrechtsen, Poul Logstrup Bjerg, Susanne Schmidt, Wei E. Huang, and Boris Maurijn Van Breukelen

Environ. Sci. Technol., **Just Accepted Manuscript** • DOI: 10.1021/acs.est.5b00715 • Publication Date (Web): 22 May 2015

Downloaded from <http://pubs.acs.org> on June 3, 2015

Just Accepted

"Just Accepted" manuscripts have been peer-reviewed and accepted for publication. They are posted online prior to technical editing, formatting for publication and author proofing. The American Chemical Society provides "Just Accepted" as a free service to the research community to expedite the dissemination of scientific material as soon as possible after acceptance. "Just Accepted" manuscripts appear in full in PDF format accompanied by an HTML abstract. "Just Accepted" manuscripts have been fully peer reviewed, but should not be considered the official version of record. They are accessible to all readers and citable by the Digital Object Identifier (DOI®). "Just Accepted" is an optional service offered to authors. Therefore, the "Just Accepted" Web site may not include all articles that will be published in the journal. After a manuscript is technically edited and formatted, it will be removed from the "Just Accepted" Web site and published as an ASAP article. Note that technical editing may introduce minor changes to the manuscript text and/or graphics which could affect content, and all legal disclaimers and ethical guidelines that apply to the journal pertain. ACS cannot be held responsible for errors or consequences arising from the use of information contained in these "Just Accepted" manuscripts.


Biodegradation: Updating the concepts of control for microbial clean-up in contaminated aquifers

Authors?

Abstract

Biodegradation is one of the most favored and sustainable means of removing organic pollutants from contaminated aquifers but the major steering factors are still surprisingly poorly understood. Growing evidence questions some of the established concepts for control of biodegradation. Here, we critically discuss classical concepts such as the thermodynamic redox zonation, or the use of steady state transport scenarios for assessing biodegradation rates. Furthermore, we discuss if absence of specific degrader populations can explain poor biodegradation. We propose updated perspectives on the controls of biodegradation in contaminant plumes. These include the plume fringe concept, transport limitations, and transient conditions as currently underestimated processes affecting biodegradation.

Abstract Art



1. Introduction

Over the last 150 years, the number of organic chemicals released into the environment has increased dramatically ¹, leaving an unprecedented chemical footprint on earth. Many groundwater contaminations result from point sources, originating from accidents or contaminations at industrial

sites. These contaminations typically form plumes with high concentrations of pollutants ($\mu\text{g/L}$ to mg/L range). Alternatively, chemicals may enter groundwater via widespread application in agriculture or release from sewage treatment into rivers. Here, pesticides, pharmaceuticals, or consumer care products are introduced as non-point sources and typically occur in much smaller concentrations (micropollutants in ng/L to $\mu\text{g/L}$ range) ².

For what seems at first sight a daunting perspective, nature fortunately has a remedy in place: biodegradation. Microorganisms can oxidize organic pollutants to CO_2 while reducing electron acceptors such as molecular oxygen, nitrate, Fe(III) (and other metal oxides), or sulfate (Fig. 1). Alternatively, some pollutants such as chlorinated solvents may serve as electron acceptors (Fig. 1, right side).




Fig. 1. Contaminants can serve as electron donors or acceptors for aquifer micro-organisms.

However, despite decades of biodegradation research, the true drivers governing contaminant degradation are still poorly understood. This article revisits and challenges current concepts on the controls and limitations of biodegradation in aquifers. It critically discusses (i) whether biodegradation is primarily governed by thermodynamics (i.e., redox zonation) at contaminated sites, (ii) if biodegradation can be adequately predicted by considering the subsurface as one reactive compartment and applying terms of environmental engineering (residence time, reaction

time), and (iii) the biological controls of biodegradation. We argue that groundwater ecosystems are much more heterogeneous and dynamic than currently perceived. Furthermore, we suggest that kinetic controls of biodegradation have been largely overlooked. Current concepts rely to a large part on thermodynamic considerations and steady state assumptions, while processes are frequently dynamic. In many cases, the steering parameters were not considered at appropriate spatial and temporal scales. However, the crucial controls of biodegradation discussed below provide potential for changing the future design of scientific projects, monitoring campaigns, or remediation strategies.

2. Revisiting redox zonation in contaminated aquifers

In highly polluted aquifers (e.g. petroleum spills with hydrocarbon concentrations of up to 100 mg/L), an excess of dissolved electron donors (hydrocarbons) prevails over acceptors (Fig. 1). In such contaminant plumes, electron acceptors are readily depleted which is widely accepted as a major limitation of biodegradation^{3,4}. A longitudinal sequence of redox processes is assumed: methanogenic degradation close to the contaminant source, followed by sulfate reduction, manganese(IV) and iron(III)-oxide reduction, nitrate reduction, and finally aerobic processes towards the distal end of the plume³⁻⁷ (Fig. 2 A). However, this redox zonation concept is challenged in the following.




Fig. 2. Comparison of the longitudinal redox zonation concept (A) and the plume fringe concept (B), both describing the spatial distribution of electron acceptors and respiration processes in a hydrocarbon contaminant plume. (B) Iron(III) reduction, manganese(IV) reduction, and methanogenesis may occur simultaneously in the core of the contaminant plume.

Dissolved electron acceptors depleted at the source zone cannot be readily replenished in the downstream plume due to laminar flow and the limited transversal dispersion in porous media (Fig. 2 B). Accordingly, methanogenic degradation or reduction of insoluble iron(III) and manganese (IV) phases would be the only electron-accepting processes possible in the source zone or the downstream plume core. Recent field evidence supports this theoretical concept showing electron acceptor depletion in the plume center^{8,9}. This is an evident contradiction to the classical concept of reverse longitudinal redox zonation (Fig. 2A). If dissolved electron acceptors such as sulfate or nitrate are consumed already at the source, they cannot become available again downstream allowing for sulfate or nitrate reduction. Even if not all electron acceptors are depleted during the passage through the source zone, a downstream redox succession should develop, where first nitrate and sulfate reduction take place, followed by methanogenesis, and not *vice versa*. Such spatial distributions have indeed been found along contaminant plumes when sampling was performed at appropriate resolution^{10,11}.

2.1. Is thermodynamics alone determining microbial competition in contaminant plumes?

The theory behind every redox zonation model is that microorganisms reducing a thermodynamically more favorable electron acceptor can gain more energy (Fig. 1), e.g. nitrate- vs. sulfate-reducing bacteria¹². In electron donor-limited systems such as aquifers with only little contamination, nitrate reducers should therefore be able to consume organic substrates to threshold concentrations no longer permissive for the activity of thermodynamically less favored respiratory guilds, which consequently become outcompeted¹³. However, in highly contaminated aquifers, electron donors are present in excess over the oxidation capacity of all electron acceptors and at concentrations much higher than where competition for electron donors (i.e., available organic substrate) can occur. Consequently, respiration processes should rather occur simultaneously as long as respective electron acceptors are present and do not become limiting for a certain respiratory guild. Biodegradation activity thus becomes controlled by availability of specific electron acceptors, rather than by thermodynamics. This concept is supported by studies on

electron acceptor-limited chemostats where axenic cultures express all respiratory pathways simultaneously rather than only the energetically most favorable¹⁴⁻¹⁷.

We propose that the reason why more favored respiratory guilds may nevertheless appear in higher abundance is a kinetic advantage which is based on higher energy conservation. Conserving more energy leads to higher growth yields (Y) and, therefore, growth rates (μ) (Equations 1-3; where X is the total biomass, X_0 the initial biomass, S the substrate concentration, and t the time of observation).

$$\mu = dX \div dt \times 1 \div X_0 \tag{1}$$

$$dX \div dt = -Y \times dS \div dt \tag{2}$$

$$\mu = -Y \times dS \div dt \times 1 \div X_0 \tag{3}$$

Thus, nitrate reducers can grow faster and to higher cell numbers in a given plume compartment suggesting an apparent out-competing of inferior respiratory guilds by thermodynamics. We propose that in many cases this will be controlled by the availability of electron acceptors and not by thermodynamic competition between respiratory guilds. Recently, Hansel et al. reported that microbial sulfate reduction was dominant over iron-reduction in sediments despite the lower thermodynamic energy gain¹⁸. The study exemplifies the importance of bioavailability rather than merely the thermodynamic redox potential of the electron acceptor.

2.2. Importance of processes at plume fringes

In recent years, it became apparent that biodegradation in contaminant plumes is much better explained by the ‘plume fringe concept’ than by the classical longitudinal redox zonation (Fig. 2)¹⁹. This concept is founded on the depletion of dissolved electron acceptors in the plume core. Biodegradation with oxygen, nitrate, or sulfate reduction can then only take place at the fringes of the plume, where electron acceptors are replenished from surrounding groundwater by dispersion and diffusion (“mixing in”) (Fig. 2 B)^{8,19-21}. At an adequately high resolution of sampling, steep geochemical counter-gradients of electron donors and acceptors have indeed been observed in several contaminated aquifers thus verifying the ‘plume fringe concept’²²⁻²⁴. The concept also provides an appropriate explanation for the overall rather limited net biodegradation rates in hydrocarbon plumes: the small-scale dispersive mixing at plume fringes controls the mass transfer of electron acceptors and, thus, microbial activities. On the other hand, the concept predicts that the plume fringes are the true hot spots where biodegradation occurs *in situ*^{25,26}.

The incomplete conceptual understanding of plume redox zonation brings about two fundamental caveats in many field investigations: (i) sampling at inappropriate scales and (ii) taking samples at the wrong spot ^{27,28}. Thus, many studies may actually have overlooked the most relevant processes and zones of biodegradation simply because groundwater sampling was done at meter rather than at centimeter resolution. While numerous studies reported on marked differences in overall microbial community assembly along plumes²⁹⁻³², vertical heterogeneity of biodegradation activities has rarely been considered ^{9,33,34}. Consequently, appropriately high resolution monitoring and the limitations of dispersive mixing still await a better incorporation into conceptual models and study design. Future research should investigate the generic conditions affecting the processes at the plume fringes and the limitations of biodegradation.

2.3. The plume core as a poorly understood compartment

Even when all dissolved electron acceptors are depleted, methanogenesis and Fe(III)- or Mn(IV)-reduction may still drive biodegradation of hydrocarbons in the plume core (Fig. 2 B) ³⁵. Evidence for iron-reduction has indeed been reported for contaminated aquifers ³⁶⁻³⁸. Very little studies, however, exist on methanogenic hydrocarbon degradation in laboratory incubations which is probably due to the extremely slow growth of such cultures ³⁹. Even less documented is methanogenic hydrocarbon degradation in aquifers ^{10,11}. Methanogenic hydrocarbon degradation may thus seem limited in spatial extent or relative contribution to the overall biodegradation ^{40,41}. An explanation might be provided by a study where agitation has been shown to impede methane production in soil slurries ⁴², most likely by disturbing close spatial interactions between syntrophic fermenters and methanogens. By analogy, groundwater flow may also interfere with efficient interspecies electron transfer in methanogenic plume zones by flushing away hydrogen or acetate and thus interfering with the energy fluxes needed for methanogenic activity ^{43,44}. Nevertheless, the true extent of methanogenic processes in contaminated aquifers requires further elucidation.

3. Bottlenecks of degradation by mass transfer




Fig. 3. Plug reactor or piston flow conceptual model of contaminated aquifers where contaminant removal (time constant of reaction, τ_{reaction}) is inversely proportional to the average residence time $\tau_{\text{transport}}$ (A). However, the distribution of flow paths on the linear pore velocity scale (B), mass transfer on the pore scale (C), mass transfer through cell membranes on the organism scale, and (bio)chemical enzymatic transformation on the molecular scale (D) can be important bottlenecks of biodegradation not taken into account by the simplified model.

3.1. Average residence times ignore heterogeneous flow paths and flow velocities

Current conceptual models frequently treat natural sediments and aquifers like either completely mixed or plug flow reactors (Fig. 3A)^{45,46}. In these models adopted from chemical engineering, a decrease in substrate concentration ($\Delta \text{conc.}$) is proportional to the residence time ($\tau_{\text{transport}}$) in the reactive compartment (Fig. 3A). The relation of transport and degradation is frequently estimated from the dimensionless Damköhler number Da

$$Da = \frac{\tau_{\text{transport}}}{\tau_{\text{reaction}}} = \frac{\text{reaction rate}}{\text{transport rate}} \quad (4)$$

where $\tau_{\text{transport}}$ is the mean residence time ("how long does it take for a compound to pass the compartment?") and τ_{reaction} the time constant of the reaction ("how long does it take for the compound to react?"). Note that the time constants are inversely correlated to the respective rates (Fig. 3A and Equation 4). The larger Da , the more biodegradation can potentially take place. This well-established concept might be a good proxy for identifying mass transfer as limiting factor in steady state systems dominated by advection. However, it is challenged by the fact that $\tau_{\text{transport}}$ and

τ_{reaction} are not well defined in natural systems because they depend on multiple parameters on different scales (Fig. 3). Black box approaches not considering multiple limitations will miss the opportunity for profound understanding of the steering parameters of such systems.

If transport is limiting on the pore velocity scale, increased flow velocities (decreasing residence times) may affect τ_{reaction} and increase biodegradation. This counterintuitive observation can be explained when considering that increased flow velocities induce more heterogeneous flow due to a wider distribution of flow paths length. This results in higher transversal and longitudinal dispersion^{47,48} and thus increases the apparent reaction rate by bringing reactants together. However, also the opposite can occur and contaminants may bypass reactive zones due to increased preferential flow. Furthermore, changes in flow velocities can create unfavorable growth conditions⁴⁸ due to shifts in nutrient fluxes and redox conditions (see section 4) adversely affecting degradation (longer τ_{reaction}). Concluding, the average residence time alone will not provide information on such ecological consequences. Rather, the distribution of residence times and the biogeochemical history along individual flow paths are governing mass transfer limitations in aquifers. This can be analyzed by high resolution monitoring which also allows for identifying dynamic flow fields and true solute fluxes⁴⁹. However, there is a need to systematically investigate the influence of flow velocities and dynamic conditions on microbial degradation.

3.2. Does diffusion limit bioavailability on the pore scale?

In groundwater, most microorganisms are attached to sediments⁵⁰ where diffusion may become the dominant mode of substrate transport to cells at the pore scale^{51,52} (Fig. 3C). If supply by diffusion is slow, biodegradation is availability-limited because microorganisms consume the substrate faster than it can be delivered⁵³. The apparent τ_{reaction} in a given environmental compartment is then determined by $\tau_{\text{diffusion}}$ (Fig. 3C). Because diffusion to the cells takes place on the micrometer scale, steep diffusive gradients can create a situation in which much larger concentrations are observed in the surrounding water. Such limitations tend to be overlooked in conventional sampling. Whether or not such diffusion limitation is important, depends also on flow velocities because diffusion gets more important if water flow velocities are low⁵⁴. This is exemplified in three scenarios regarding concentrations and the state of the system.

(i) At *high concentrations and steady state*, diffusive gradients between pore centers and sediment surfaces only build up if water flow velocities are low and if degradation rates are faster than the supply of contaminants. (ii) For *low concentrations and transient conditions* Langner *et al.* found that degradation rates of 2,4-dichlorophenoxyacetic acid were higher when water flow was slower, even

though the flow path was shorter⁵⁵. This indicates that molecules needed sufficient time to diffuse into micropores and $\tau_{\text{transport}}$ has to be sufficiently long compared to $\tau_{\text{diffusion}}$, since otherwise molecules were flushed through the pores without degradation. (iii) For *low to medium concentrations and at steady state*, one can expect that diffusive gradients become shallow if $\tau_{\text{transport}}$ is long and concentrations are low. Consequently, also the substrate supply by diffusive fluxes becomes slower according to Fick's law. Then, higher flow velocities would actually be advantageous because they replenish the substrate creating steeper gradients and increasing diffusive substrate supply to the organisms. Thus, the pore water velocity is an important parameter contributing to diffusion limitations on the pore scale. However, both, too high and too small pore water velocities can induce limitations in bioavailability⁵¹ which implies a need for a systematic elucidation of this topic.

3.3. Thermodynamics, mass transfer, or enzyme kinetics: what is limiting on the organism scale?

It is often observed that micropollutants are only incompletely degraded even when competent bacterial degraders are present⁵⁶. This unsolved paradox of threshold concentrations might be due to different reasons. (i) Thermodynamic limitation for biodegradation is an often considered explanation, but can typically be excluded. For example at nM toluene concentrations, the Gibbs enthalpy of reaction ΔG for aerobic degradation (-3890 kJ/mol) would be large enough to consume even the last toluene molecule. At high dilution, biodegradation is rather kinetically limited by mass transfer to the cell as explained above. (ii) An alternative explanation for incomplete degradation is a kinetic limitation by insufficient substrate uptake into the cell⁵⁷. (iii) Furthermore, slow biochemical transformation rates (enzyme kinetics) might be due to the intrinsically difficult-to-degrade molecular structures of xenobiotics⁵⁸. This is supported by comparably slow degradation rates of persistent compounds at higher but non-toxic concentrations in batch cultures⁵⁹.

In natural systems, it was so far not possible to distinguish the different kinds of limitation of biodegradation on the organism scale, which opens future research fields.

4. Microbial controls of biodegradation

Absence of specific degrader populations is often assumed when insufficient biodegradation is observed at a given site. At organohalide-contaminated sites, bioaugmentation (amending respective degraders) of microbial consortia containing e.g. *Dehalococcoides* (Dhc) strains capable of reductive dechlorination of trichloroethylene (TCE) to ethene has been successful⁶⁰⁻⁶². Similarly, the effectiveness of bioaugmentation in atrazine- and MTBE-contaminated (methyl-tert-butylether) aquifers has been demonstrated⁶³⁻⁶⁵. However, even highly specialized organohalide-respiring

bacteria are generally widespread in aquifers⁶⁶⁻⁶⁸. Thus for certain settings, it remains questionable whether respective degrader organisms were truly absent before bioaugmentation or only present at very low abundance.

4.1. Limitation of biodegradation by microbial growth.

In aquifers, a substantial fraction of the microbes is suggested to be in a status of low activity, inactive, or even dormant⁶⁹. Moreover, microbial communities in aquatic systems exhibit growth efficiencies (yields) much lower than known from batch cultures and chemostats, with median values below 0.3 (g biomass / g substrate) for rivers, lakes, and oceans^{70,71}. *In situ* growth rates are also low (equation 3) for aquifer systems, and doubling times can be in the range of months to years⁷². Under optimum conditions in the laboratory, the presence of one degrader cell at the moment of a hydrocarbon spill would allow aerobes to form notable biomass (e.g. 10^5 to 10^6 cells per liter groundwater) within a day, while e.g. sulfate reducers or organohalide reducers (doubling time: ~ 10 d) may require >100 d to establish a critical population size. Indeed, a fast response has been observed for an oxic aquifer system receiving a short contaminant pulse⁷³, while anaerobic degradation coupled to denitrification established only over several weeks. For more recalcitrant compounds and pollutants, requiring anoxic conditions for degradation (such as halogenated solvents), it might take years before reasonable numbers of degraders have developed. Thus in aquifers, a slow community response might be misinterpreted as absence of degrader populations which needs to be verified in the future.

4.2. Limitations of biodegradation by microbial physiology

Total concentrations of dissolved organic carbon (DOC) in pristine groundwater are usually in the low mg L⁻¹ range (0.5-2 mg L⁻¹), of which only 0.5 to 5% are readily assimilable organic carbon (AOC)⁷⁴⁻⁷⁶. This AOC consists of a plethora of individual compounds at extremely low individual concentrations, including organic micropollutants such as pesticides, pharmaceuticals, and many other low-level contaminants⁷⁷. The latter are often present below the threshold concentrations of initial induction of catabolic genes and degradation pathways⁷⁵. At very low concentrations in the environment, it is likely that microorganisms do not feed on only one substrate at a time. Mixed substrate utilization – where microbes can utilize a wide range of offered substrates simultaneously – has been observed in carbon-limited chemostats⁷⁸. This leads to much lower threshold concentrations for individual compounds implying that the degradation of one compound “helps” to degrade another compound in energetic co-metabolism. However, at excess of substrate in hydrocarbon plumes, catabolite repression, competitive inhibition, or metabolic flux dilution might take place^{79,80}.

Today, it is totally unclear how microbial metabolism is regulated in the environment. Such knowledge will be useful for designing strategies for removing micropollutants from drinking water in engineered systems or improving the licensing practice of pesticides⁷⁷.

4.3. Limitations of ecosystem response

Low microbial growth rates imply that even under steady state conditions where organisms can develop without disturbance, long time spans are required to establish significant degrader biomass and thus biodegradation capacities. However in contaminated groundwater, conditions are not necessarily in steady state⁸¹⁻⁸³. In fact, temporal hydraulic fluctuations may represent a major control of biodegradation in groundwater by repeatedly changing the environmental conditions encountered by degraders (e.g. sudden exposure of anaerobes to oxygen). This can be exemplified by plume fringes which are characterized not only as hot spots of biodegradation activity, but also by an apparently 'specialized' degrader microbiota^{25,9} over only a few dm. If the prevailing geochemical conditions for microorganisms shift, degraders must continuously follow or re-establish in other strata. Geochemical shifts of the plume could thus represent a further, as-yet unrecognized kinetic limitation of biodegradation.

Moreover, transient supply of substrates by such fluctuations may not be sufficient to support growth of degrader populations. It can be speculated that under spatially and temporarily dynamic hydraulic conditions, degrader populations may never reach the biomass levels required to efficiently degrade substrate pulses. Once formed, however, degrader biomass may persist for months and perhaps even years after the source of contamination has disappeared^{73,84}. Biomass established upon previous locations of the plume could sustain biodegradation capacities for future contaminations. Thus, aquifers could become preconditioned to efficiently degrade pollutants.

4.4. Further research needs.

The role of grazers (protozoa) and viruses (phages) in shaping prokaryotic degrader communities and influencing *in situ* degradation rates is totally overlooked to date⁸⁵. While the influence of protozoa on bacterial community composition and *vice versa* has been shown also for contaminated groundwater habitats^{86,87}, there is contradicting evidence on either the stimulation^{88,89} or inhibition^{90,91} of biodegradation by protozoan grazing. Only a few studies are available on bacteriophages in groundwater⁹²⁻⁹⁴ but their influence on degrader communities and activities have not been addressed, so far. Extrapolating recent advances from surface aquatic environments and marine systems, bacteriophages can be expected to play a significant role in controlling prokaryotic

production and diversity. With reference to the highly specialized degrader populations found in biodegradation hot spots, ecological concepts such as the ‘killing the winner’ hypothesis await to be tested. It predicts that when a given population grows beyond a critical density, grazers and viruses will decimate the population⁹⁵ affecting biodegradation.

5. Conclusion and outlook

Here we discuss several controls for biodegradation in contaminated aquifers that have been recognized in recent years, and call for an update of classical black box approaches in site assessment and restoration. New perspectives in groundwater research should include the plume fringe concept and mass transfer limitations as steering factors for biodegradation. On the organism scale, physiological properties of degraders and ecological drivers of degrader community structure have been identified to affect biodegradation. We propose that biodegradation in contaminated aquifers is largely controlled by kinetics. Different kinetic controls are interacting in complex ways and cannot be described by flow- or residence-time-dependent degradation rates alone. An important caveat is that many of these mechanisms act at the μm -to-cm scale, while sampling is still mostly conducted at the meter scale. To fully understand process limitations, samples have to be taken at adequate resolution, often including intact sediment cores or highly-resolved water sampling. Furthermore, temporal dynamics of processes demand for extended monitoring with more frequent sampling intervals in time and space.

Acknowledgement: This work was funded by the EU-initial training network “Goodwater” and the Helmholtz Zentrum München.

6. References

1. Schwarzenbach, R.; Egli, T.; Hofstetter, T. B.; von Gunten, U.; Wehrli, B., Global water pollution and human health. *Ann. Rev. Environ. Res.* **2010**, *35*, 109-136.
2. Richardson, S. D.; Ternes, T. A., Water analysis: emerging contaminants and current issues. *Anal.Chem.* **2011**, *83*, 4614-4648.
3. Christensen, T. H.; Kjeldsen, P.; Albrechtsen, H.-J.; Heron, G.; Nielsen, P. H.; Bjerg, P. L.; Holm, P. E., Attenuation of landfill leachate pollutants in aquifers. *Crit. Rev. Environ. Sci. Technol.* **1994**, *24*, 119-202.

- 332 4. Wiedemeier, T. H.; Rifai, H. S.; Newell, C. J.; Wilson, J. T., *Natural attenuation of fuels and*
333 *chlorinated solvents in the subsurface*. Wiley: New York, 1999.
- 334 5. Chapelle, F. H., The significance of microbial processes in hydrogeology and geochemistry.
335 *Hydrogeol. J.* **2000**, *8*, 41-46.
- 336 6. Lovley, D. R., Cleaning up with genomics: Applying molecular biology to bioremediation.
337 *Nature Rev. Microbiol.* **2003**, *1*, 35-44.
- 338 7. Christensen, T. H.; Bjerg, P. L.; Banwart, S. A.; Jakobsen, R.; Heron, G.; Albrechtsen, H. J.,
339 Characterization of redox conditions in groundwater contaminant plumes. *Journal of Contaminant*
340 *Hydrology* **2000**, *45*, 165-241.
- 341 8. Anneser, B.; Einsiedl, F.; Meckenstock, R. U.; Richters, L.; Wisotzky, F.; Griebler, C., High-
342 resolution monitoring of biogeochemical gradients in a tar oil-contaminated aquifer. *Appl. Geochem.*
343 **2008**, *23*, 1715-1730.
- 344 9. Anneser, B.; Pilloni, G.; Bayer, A.; Lueders, T.; Griebler, C.; Einsiedl, F.; Richters, L., High
345 resolution analysis of contaminated aquifer sediments and groundwater—what can be learned in
346 terms of natural attenuation? *Geomicrobiol. J.* **2010**, *27*, 130 - 142.
- 347 10. Tischer, K.; Kleinstaub, S.; Schleinitz, K. M.; Fetzer, I.; Spott, O.; Stange, F.; Lohse, U.; Franz,
348 J.; Neumann, F.; Gerling, S.; Schmidt, C.; Hasselwander, E.; Harms, H.; Wendeberg, A., Microbial
349 communities along biogeochemical gradients in a hydrocarbon-contaminated aquifer. *Environ.*
350 *Microbiol.* **2013**, *15*, 2603-2615.
- 351 11. Fahrenfeld, N.; Cozzarelli, I.; Bailey, Z.; Pruden, A., Insights into biodegradation through
352 depth-resolved microbial community functional and structural profiling of a crude-oil contaminant
353 plume. *Microb. Ecol.* **2014**, *68*, 453-462.
- 354 12. Zehnder, A. J. B.; Stumm, W., Geochemistry and biochemistry of anaerobic habitats. In
355 *Biology of Anaerobic Microorganisms*, Zehnder, A. J. B., Ed. John Wiley & Sons: New York, 1988; pp
356 1-38.
- 357 13. Achtnich, C.; Bak, F.; Conrad, R., Competition for electron donors among nitrate reducers,
358 ferric iron reducers, sulfate reducers, and methanogens in anoxic paddy soil. *Biol. Fertil. Soils* **1995**,
359 *19*, 65-72.
- 360 14. Ihssen, J.; Egli, T., Global physiological analysis of carbon- and energy-limited growing
361 *Escherichia coli* confirms a high degree of catabolic flexibility and preparedness for mixed substrate
362 utilization. *Environmental Microbiology* **2005**, *7*, 1568-1581.
- 363 15. Kovarova-Kovar, K.; Egli, T., Growth kinetics of suspended microbial cells: From single-
364 substrate-controlled growth to mixed-substrate kinetics. *Microbiol. Mol. Biol. Rev.* **1998**, *62*, 646-+.
- 365 16. Marozava, S.; Röling, W. F. M.; Seifert, J.; Küffner, R.; von Bergen, M.; Meckenstock, R. U.,
366 Physiology of *Geobacter metallireducens* under excess and limitation of electron donors. Part I.
367 Batch cultivation with excess of carbon sources. *Syst. Appl. Microbiol.* **2014**, *37*, 277-286.
- 368 17. Marozava, S.; Röling, W. F. M.; Seifert, J.; Küffner, R.; von Bergen, M.; Meckenstock, R. U.,
369 Physiology of *Geobacter metallireducens* under excess and limitation of electron donors. Part II.

- 370 Mimicking environmental conditions during cultivation in retentostats. *Syst. Appl. Microbiol.* **2014**,
371 37, 287-295.
- 372 18. Hansel, C. M.; Lentini, C. J.; Tang, Y.; Johnston, D. T.; Wankel, S. D.; Jardine, P. M., Dominance
373 of sulfur-fueled iron oxide reduction in low-sulfate freshwater sediments. *ISME J* **2015**,
374 (10.1038/ismej.2015.50).
- 375 19. Bauer, R. D.; Maloszewski, P.; Zhang, Y. C.; Meckenstock, R. U.; Griebler, C., Mixing-
376 controlled biodegradation in a toluene plume - Results from two-dimensional laboratory
377 experiments. *J. Cont. Hydrol.* **2008**, 96, 150-168.
- 378 20. Cirpka, O. A.; Frind, E. O.; Helmig, R., Numerical simulation of biodegradation controlled by
379 transverse mixing. *J. Contam. Hydrol.* **1999**, 40, 159-182.
- 380 21. Chu, M.; Kitanidis, P. K.; McCarty, P. L., Modeling microbial reactions at the plume fringe
381 subject to transverse mixing in porous media: When can the rates of microbial reaction be assumed
382 to be instantaneous? *Water Res. Res.* **2005**, 41, 1-15.
- 383 22. Thornton, S.; Quigley, S.; Spence, N. J.; Banwart, S.; Bottrell, S.; Lerner, D. N., Processes
384 controlling the distribution and natural attenuation of dissolved phenolic compounds in a deep
385 sandstone aquifer. *J. Contam. Hydrol.* **2001**, 53, 233-267.
- 386 23. Van Breukelen, B. M.; Griffioen, J., Biogeochemical processes at the fringe of a landfill
387 leachate pollution plume: potential for dissolved organic carbon, Fe(II), Mn(II), NH₄, and CH₄
388 oxidation. *J. Contam. Hydrol.* **2004**, 73, 181-205.
- 389 24. Tuxen, N.; Albrechtsen, H.-J.; Bjerg, P. L., Identification of a reactive degradation zone at a
390 landfill leachate plume fringe using high resolution sampling and incubation techniques. *J. Contam.*
391 *Hydrol.* **2006**, 85, 179-194.
- 392 25. Winderl, C.; Anneser, B.; Griebler, C.; Meckenstock, R. U.; Lueders, T., Depth-Resolved
393 Quantification of Anaerobic Toluene Degradation and Aquifer Microbial Community Patterns in
394 Distinct Redox Zones of a Tar Oil Contaminant Plume. *Appl. Environ. Microbiol.* **2008**, 74, 792-801.
- 395 26. Jobelius, C.; Ruth, B.; Griebler, C.; Meckenstock, R. U.; Hollender, J.; Reineke, A.; Frimmel, F.
396 H.; Zwiener, C., Metabolites indicate hot spots of biodegradation and biogeochemical gradients in a
397 high-resolution monitoring well. *Environ. Sci. Technol.* **2010**, 45, 474-481.
- 398 27. Lehman, R. M., Understanding of aquifer microbiology is tightly linked to sampling
399 approaches. *Geomicrobiol. J.* **2007**, 24, 331-341.
- 400 28. Wilson, R. D.; Thornton, S. F.; Mackay, D. M., Challenges in monitoring the natural
401 attenuation of spatially variable plumes. *Biodegradation* **2004**, 15, 359-369.
- 402 29. Dojka, M. A.; Hugenholtz, P.; Haack, S. K.; Pace, N. R., Microbial diversity in a hydrocarbon-
403 and chlorinated-solvent- contaminated aquifer undergoing intrinsic bioremediation. *Appl. Environ.*
404 *Microbiol.* **1998**, 64, 3869-3877.
- 405 30. Feris, K. P.; Hristova, K.; Gebreyesus, B.; Mackay, D.; Scow, K. M., A shallow BTEX and MTBE
406 contaminated aquifer supports a diverse microbial community. *Microb. Ecol.* **2004**, 48, 589-600.

- 407 31. Yagi, J. M.; Neuhauser, E. F.; Ripp, J. A.; Mauro, D. M.; Madsen, E. L., Subsurface ecosystem
408 resilience: long-term attenuation of subsurface contaminants supports a dynamic microbial
409 community. *ISME J.* **2010**, *4*, 131-143.
- 410 32. Staats, M.; Braster, M.; Roling, W. F. M., Molecular diversity and distribution of aromatic
411 hydrocarbon-degrading anaerobes across a landfill leachate plume. *Environ. Microbiol.* **2011**, *13*,
412 1216-1227.
- 413 33. Winderl, C.; Anneser, B.; Griebler, C.; Meckenstock, R. U.; Lueders, T., Depth-resolved
414 quantification of anaerobic toluene degraders and aquifer microbial community patterns in distinct
415 redox zones of a tar oil contaminant plume. *Appl. Environ. Microbiol.* **2008**, *74*, 792-801.
- 416 34. Tischer, K.; Kleinstaub, S.; Schleinitz, K. M.; Fetzer, I.; Spott, O.; Stange, F.; Lohse, U.; Franz,
417 J.; Neumann, F.; Gerling, S.; Schmidt, C.; Hasselwander, E.; Harms, H.; Wendeberg, A., Microbial
418 communities along biogeochemical gradients in a hydrocarbon-contaminated aquifer. *Environ.*
419 *Microbiol.* **2013**, *15*, 2603-2615.
- 420 35. Foght, J., Anaerobic biodegradation of aromatic hydrocarbons: Pathways and prospects. *J.*
421 *Mol. Microbiol. Biotechnol.* **2008**, *15*, 93-120.
- 422 36. Christensen, T. H.; Kjeldsen, P.; Albrechtsen, H.-J.; Heron, G.; Nielsen, P. H.; Bjerg, P. L.; Holm,
423 P. E., Attenuation of landfill leachate pollutants in aquifers. *Crit. Rev. Environ. Sci. Technol.* **1994**, *24*,
424 119-202.
- 425 37. Albrechtsen, H.-J.; Christensen, T., Evidence for microbial iron reduction in a landfill
426 leachate-polluted aquifer (Vejen, Denmark). *Appl. Environ. Microbiol.* **1994**, *60*, 3920-3925.
- 427 38. Botton, S.; Van Harmelen, M.; Braster, M.; Parsons, J. R.; Röling, W. F. M., Dominance of
428 *Geobacteraceae* in BTX-degrading enrichments from an iron-reducing aquifer. *FEMS Microbiol. Ecol.*
429 **2007**, *62*, 118-130.
- 430 39. Gieg, L. M.; Fowler, S. J.; Berdugo-Clavijo, C., Syntrophic biodegradation of hydrocarbon
431 contaminants. *Curr. Op. Biotechnol.* **2014**, *27*, 21-19.
- 432 40. Bekins, B. A.; Cozzarelli, I. M.; Godsy, E. M.; Warren, E.; Essaid, H. I.; Tuccillo, M. E.,
433 Progression of natural attenuation processes at a crude oil spill site: II. Controls on spatial
434 distribution of microbial populations. *J. Contam. Hydrol.* **2001**, *53*, 387-406.
- 435 41. Feisthauer, S.; Seidel, M.; Bombach, P.; Traube, S.; Knöller, K.; Wange, M.; Fachmann, S.;
436 Richnow, H. H., Characterization of the relationship between microbial degradation processes at a
437 hydrocarbon contaminated site using isotopic methods. *J. Contam. Hydrol.* **2012**, *133*, 17-29.
- 438 42. Dannenberg, S.; Wudler, J.; Conrad, R., Agitation of anoxic paddy soil slurries affects the
439 performance of the methanogenic microbial community. *FEMS Microbiol. Ecol.* **1997**, *22*, 257-263.
- 440 43. Schink, B., Synergistic interactions in the microbial world. *Antonie van Leeuwenhoek* **2002**,
441 *81*, 257-261.
- 442 44. Stams, A. J. M.; Plugge, C. M., Electron transfer in syntrophic communities of anaerobic
443 bacteria and archaea. *Nature Rev. Microbiol.* **2009**, *7*, 568-577.

- 444 45. Carleton, J. N., Damköhler number distributions and constituent removal in treatment
445 wetlands. *Ecol. Engin.* **2002**, *19*, 233-248.
- 446 46. Oldham, C. E.; Farrow, D. E.; Peiffer, S., A generalized Damköhler number for classifying
447 material processing in hydrological systems. *Hydrol. Earth Syst. Sci.* **2013**, *17*, 1133-1148.
- 448 47. Song, X.; Seagren, E. A., In Situ Bioremediation in Heterogeneous Porous Media: Dispersion-
449 Limited Scenario. *Environ. Sci. Technol.* **2008**, *42*, 6131-6140.
- 450 48. Mendoza-Sanchez, I.; Autenrieth, R. L.; McDonald, T. J.; Cunningham, J. A., Effect of pore
451 velocity on biodegradation of *cis*-dichloroethene (DCE) in column experiments. *Biodegradation* **2010**,
452 *21*, 365-377.
- 453 49. Rein, A.; Bauer, S.; Dietrich, P.; al., e., Influence of temporally variable groundwater flow
454 conditions on point measurements and contaminant mass flux estimations *J. Cont. Hydrol.* **2009**,
455 *108*, 118-133.
- 456 50. Griebler, C.; Lueders, T., Microbial biodiversity in groundwater ecosystems. *Freshwater Biol.*
457 **2009**, *54*, 649-677.
- 458 51. Hesse, F.; Harms, H.; Attinger, S.; Thullner, M., Linear exchange model for the description of
459 mass transfer limited bioavailability at the pore scale. *Environ. Sci. Technol.* **2010**, *44*, 2064-2071.
- 460 52. Ramaswami, A.; Ghoshal, S.; Luthy, R. G., Mass transfer and bioavailability of PAH
461 compounds in coal tar NAPL-slurry systems. 2. Experimental evaluations. *Environ. Sci. Technol.* **1997**,
462 *31*, 2268-2276.
- 463 53. Bosma, T. N. P.; Middeldorp, P. J. M.; Schraa, G.; Zehnder, A. J. B., Mass transfer limitation of
464 biotransformation: quantifying bioavailability. *Environ. Sci. Technol.* **1997**, *31*, 248-252.
- 465 54. Brusseau, M. L., The influence of solute size, pore water velocity, and intraparticle porosity
466 on solute dispersion and transport in soil. *Water Res. Res.* **1993**, *29*, 1071-1080.
- 467 55. Langner, H. W.; Inskeep, W. P.; Gaber, H. M.; Jones, W. L.; Das, B. S.; Wraith, J. M., Pore
468 water velocity and residence time effects on the degradation of 2,4-D during transport. *Environ. Sci.*
469 *Technol.* **1998**, *32*, 1308-1315.
- 470 56. Janniche, G. S.; Mouvet, C.; Albrechtsen, H. J., Vertical small scale variations of sorption and
471 mineralization of three herbicides in subsurface limestone and sandy aquifer. *J. Contam. Hydrol.*
472 **2011**, *123*, 167-177.
- 473 57. Ferenci, T.; Robert, K. P., Bacterial physiology, regulation and mutational adaptation in a
474 chemostat environment. *Adv. Microb. Physiol.* **2008**, *53*, 169-229.
- 475 58. Rücker, C.; Kümmerer, K., Modeling and predicting aquatic aerobic biodegradation - a review
476 from a user's perspective. *Green Chem.* **2012**, *14*, 875-887.
- 477 59. Liu, J.; Mejia Avendaño, S., Microbial degradation of polyfluoroalkyl chemicals in the
478 environment: A review. *Environ. Internat.* **2013**, *61*, 98-114.
- 479 60. Paliwal, V.; Puranik, S.; Purohit, H. J., Integrated perspective for effective bioremediation.
480 *Appl. Biochem. Biotechnol.* **2012**, *166*, 903-924.


- 481 61. Tyagi, M.; da Fonseca, M.; de Carvalho, C., Bioaugmentation and biostimulation strategies to
482 improve the effectiveness of bioremediation processes. *Biodegradation* **2011**, *22*, 231-241.
- 483 62. Major, D. W.; McMaster, M. L.; Cox, E. E.; Edwards, E. A.; Dworatzek, S. M.; Hendrickson, E.
484 R.; Starr, M. G.; Payne, J. A.; Buonamici, L. W., Field demonstration of successful bioaugmentation to
485 achieve dechlorination of tetrachloroethene to ethene. *Environ. Sci. Technol.* **2002**, *36*, 5106-5116.
- 486 63. Salanitro, J. P.; Johnson, P. C.; Spinnler, G. E.; Maner, P. M.; Wisniewski, H. L.; Bruce, C., Field-
487 scale demonstration of enhanced MTBE bioremediation through aquifer bioaugmentation and
488 oxygenation. *Environ. Sci. Technol.* **2000**, *34*, 4152-4162.
- 489 64. Shati, M. R.; Rönen, D.; Mandelbaum, R., Method for *in situ* study of bacterial activity in
490 aquifers. *Environ. Sci. Technol.* **1996**, *30*, 2646-2653.
- 491 65. Smith, A. E.; Hristova, K.; Wood, I.; Mackay, D. M.; Lory, E.; Lorenzana, D.; Scow, K. M.,
492 Comparison of biostimulation versus bioaugmentation with bacterial strain PM1 for treatment of
493 groundwater contaminated with methyl tertiary butyl ether (MTBE). *Environ. Health Persp.* **2005**,
494 *113*, 317-322.
- 495 66. Hendrickson, E.; Payne, J.; Young, R.; Starr, M.; Perry, M.; Fahnestock, S.; Ellis, D. E.; Ebersole,
496 R., Molecular analysis of *Dehalococcoides* 16S ribosomal DNA from chloroethene-contaminated sites
497 throughout North America and Europe. *Appl. Environ. Microbiol.* **2002**, *68*, 485-495.
- 498 67. Regeard, C.; Maillard, J.; Dufraigne, C.; Deschavanne, P.; Holliger, C., Indications for
499 acquisition of reductive dehalogenase genes through horizontal gene transfer by *Dehalococcoides*
500 *ethenogenes* strain 195. *Appl. Environ. Microbiol.* **2005**, *71*, 2955-2961.
- 501 68. Scheutz, C.; Durant, N. d.; Dennis, P.; Hansen, M. H.; Jørgensen, T.; Jakobsen, R.; Cox, E. e.;
502 Bjerg, P. L., Concurrent ethene generation and growth of *Dehalococcoides* containing vinyl chloride
503 reductive dehalogenase genes during an enhanced reductive dechlorination field demonstration.
504 *Environ. Sci. Technol.* **2008**, *42*, 9302-9309.
- 505 69. Kieft, T. L.; Phelps, T. J., Life in the slow lane: activities of microorganisms in the subsurface.
506 In *The microbiology of the terrestrial deep subsurface*, Amy, P. S.; Haldeman, D. L., Eds. Lewis
507 Publishers: Boca Raton, 1997; pp 137-163.
- 508 70. del Giorgio, P.; Cole, J., Bacterial growth efficiency in natural aquatic systems. *Ann. Rev. Ecol.*
509 *Syst.* **1998**, *29*, 503-541.
- 510 71. Sinsabaugh, R. L.; Manzoni, S.; Moorhead, D. L.; Richter, A., Carbon use efficiency of
511 microbial communities: stoichiometry, methodology and modelling. *Ecol. Lett.* **2013**, *16*, 930-939.
- 512 72. Albrechtsen, H. J.; Winding, A., Microbial biomass and activity in subsurface sediment from
513 Vejen, Denmark. *Microb. Ecol.* **1992**, *23*, 303-317.
- 514 73. Herzyk, A.; Maloszewski, P.; Qiu, S.; Elsner, M.; Griebl, C., Intrinsic potential for immediate
515 biodegradation of toluene in a pristine, energy-limited aquifer. *Biodegradation* **2014**, *25*, 325-336.
- 516 74. Goody, D. C.; Hinsby, K., Organic quality of groundwaters. In *Natural groundwater quality*,
517 Edmunds, W. M.; Shand, P., Eds. Blackwell Publishing: Oxford, 2008; pp 59-70.
- 518 75. Egli, T., How to live at very low substrate concentration. *Water Res.* **2010**, *44*, 4826-4837.

- 519 76. Zhou, Y.; Kellermann, C.; Griebler, C., Spatio-temporal patterns of microbial communities in a
520 hydrologically dynamic pristine aquifer. *FEMS Microbiol. Ecol.* **2012**, *81*, 230-242.
- 521 77. Benner, J.; Helbling, D. E.; Kohler, H.-P. E.; Wittebol, J.; Kaiser, E.; Prasse, C.; Ternes, T. A.;
522 Albers, C. N.; Aamand, J.; Horemans, B.; Springael, D.; Walravens, E.; Boon, N., Is biological
523 treatment a viable alternative for micropollutant removal in drinking water treatment processes?
524 *Water Res.* **2013**, *47*, 5955-5976.
- 525 78. Egli, T., The ecological and physiological significance of the growth of heterotrophic
526 microorganisms with mixtures of substrates. *Adv. Microb. Ecol.* **1995**, *14*, 305-386.
- 527 79. Meckenstock, R. U.; Warthmann, R. I.; Schafer, W., Inhibition of anaerobic microbial o-xylene
528 degradation by toluene in sulfidogenic sediment columns and pure cultures. *FEMS Microbiol. Ecol.*
529 **2004**, *47*, 381-386.
- 530 80. Lovanh, N.; Alvarez, P. J. J., Effect of ethanol, acetate, and phenol on toluene degradation
531 activity and tod-lux expression in *Pseudomonas putida* TOD102: Evaluation of the metabolic flux
532 dilution model. *Biotechnol. Bioeng.* **2004**, *86*, 801-808.
- 533 81. Fretwell, B. A.; Burgess, W. G.; Barker, J. A.; Jefferies, N. L., Redistribution of contaminants by
534 a fluctuating water table in a micro-porous, double-porosity aquifer: Field observations and model
535 simulations. *J. Contam. Hydrol.* **2005**, *78*, 27-52.
- 536 82. Scholl, M. A.; Cozzarelli, I. M.; Christenson, S. C., Recharge processes drive sulfate reduction
537 in an alluvial aquifer contaminated with landfill leachate. *J. Contam. Hydrol.* **2006**, *86*, 239-261.
- 538 83. van Stempvoort, D. R.; Armstrong, J.; Mayer, B., Seasonal recharge and replenishment of
539 sulfate associated with biodegradation of a hydrocarbon plume. *Groundwater Monitor. Remed.*
540 **2007**, *27*, 110-121.
- 541 84. Mellage, A.; Eckert, D.; Grösbacher, M.; Griebler, C.; Cirpka, O. A., Microbial dynamics during
542 contaminant degradation in aerobic flow-through systems under growth and starvation conditions.
543 *Environ. Sci. Technol.* **2015**, *submitted*.
- 544 85. Griebler, C.; Malard, F.; Lefébure, T., Current developments in groundwater ecology — from
545 biodiversity to ecosystem function and services. *Curr. Op. Biotechnol.* **2014**, *27*, 159-167.
- 546 86. Holmes, D.; Giloteaux, L.; Williams, K.; Wrighton, K.; Wilkins, M.; Thomson, C.; Roper, T.;
547 Long, P.; Lovley, D., Enrichment of specific protozoan populations during *in situ* bioremediation of
548 uranium-contaminated groundwater. *ISME J.* **2013**, *7*, 1286-1298.
- 549 87. Madsen, E.; Sinclair, J.; Ghiorse, W., In situ biodegradation: microbiological patterns in a
550 contaminated aquifer. *Science* **1991**, *252*, 830-833.
- 551 88. Mattison, R.; Taki, H.; Harayama, S., The soil flagellate *Heteromita globosa* accelerates
552 bacterial degradation of alkylbenzenes through grazing and acetate excretion in batch culture.
553 *Microb. Ecol.* **2005**, *49*, 142-150.
- 554 89. Tso, S.-F.; Taghon, G., Protozoan grazing increases mineralization of naphthalene in marine
555 sediment. *Microb. Ecol.* **2006**, *51*, (4), 460-469.


- 556 90. Kota, S.; Borden, R.; Barlaz, M., Influence of protozoan grazing on contaminant
557 biodegradation. *FEMS Microbiol. Ecol.* **1999**, *29*, 179-189.
- 558 91. Cunningham, J. J., III; Kinner, N. E.; Lewis, M., Protistan predation affects trichloroethene
559 biodegradation in a bedrock aquifer. *Appl. Environ. Microbiol.* **2009**, *75*, (24), 7588-7593.
- 560 92. Smith, R. J.; Jeffries, T. C.; Roudney, B.; Seymour, J. R.; Fitch, A. J.; Simons, K. L.; Speck, P. G.;
561 Newton, K.; Brown, M. H.; Mitchell, J. G., Confined aquifers as viral reservoirs. *Environ. Microbiol.*
562 *Rep.* **2013**, *5*, 725-730.
- 563 93. Eydal, H. S. C.; Jägevall, S.; Hermansson, M.; Pedersen, K., Bacteriophage lytic to
564 *Desulfovibrio aespoeensis* isolated from deep groundwater. *ISME J.* **2009**, *3*, 1139-1147.
- 565 94. Kyle, J. E.; Eydal, H. S. C.; Ferris, F. G.; Pedersen, K., Viruses in granitic groundwater from 69
566 to 450 m depth of the Äspö Hard Rock Laboratory, Sweden. *ISME J.* **2008**, *2*, 571-574.
- 567 95. Winter, C.; Bouvier, T.; Weinbauer, M.; Thingstad, T., Trade-offs between competition and
568 defense specialists among unicellular planktonic organisms: the 'killing the winner' hypothesis
569 revisited. *Microbiol. Mol. Biol. Rev.* **2010**, *74*, (1), 42-57.

570

ELECTRON DONORS




Natural Organic Matter (NOM)

<CH₂O>


ELECTRON ACCEPTORS FOR RESPIRATION

Natural Electron Acceptors





Reducible Pollutants

CHLORINATED HYDROCARBONS





NITROAROMATICS






(A) Current Conceptual Models





(B) Heterogeneous Flow Paths on the Linear Pore Velocity Scale



(C) Mass Transfer on the Pore Scale ...

(D) ... and on the Organism Scale





85x47mm (300 x 300 DPI)